

## XIX CONGRESO DE LA SOCIEDAD ARGENTINA DE MICROBIOLOGÍA GENERAL

22 al 25 de octubre del 2024 Centro cultural y Pabellón Argentina de la Universidad Nacional de Córdoba, Córdoba, ARGENTINA.



Foto: Se hace camino al andar. Celeste Dea. 1er puesto. Concurso fotográfico SAMIGE 20 años.

## TAM SYSTEM IS INVOLVED IN PHOSPHOLIPID HOMEOSTASIS IN THE OPPORTUNISTIC PATHOGEN Ochrobactrum anthropi

Zimmerman Luciana M., Russo Daniela M, Poschenrieder Julieta, Pellizza Leonardo, Aran Martín, Zorreguieta Angeles, Bialer Magalí G.

Fundación Instituto Leloir, IIBBA-CONICET - CABA - Buenos Aires - Argentina *Contacto: magalibialer@gmail.com* 

The precise assembly and remodeling of cell envelope structures in Gramnegative bacteria, especially the outer membrane (OM), is crucial for successful infection and resistance to diverse environments. The OM consists of an asymmetric lipid bilayer composed of lipopolysaccharide (LPS) on the outer leaflet and phospholipids (PLs) on the inner leaflet. The OM contains proteins (OMPs) with a ?-barrel structure and lipoproteins, some of which interact with the peptidoglycan. Despite considerable advances in understanding the mechanisms underlying OM biogenesis, certain processes are still unknown. Evidence was presented indicating that in gammaproteobacteria, the TAM system would be involved in the correct translocation of a subset of OMPs. TAM consists of TamA, a member of the Omp85 superfamily, and TamB, a large protein of the AsmA-like family that is inserted in the inner membrane (IM) by an N-terminal non-cleavable signal peptide while the rest of the protein is immersed in the periplasm. Brucella is an intracellular pathogen from the Alphaproteobacteria group that exhibits unique cell surface characteristics, making it highly resistant to various disruptive agents. While we found that certain OMPs may not be efficiently translocated to the OM in a mutant of the TamB homologue (MapB) of Brucella suis, several pieces of evidence indicate that MapB plays a crucial role in OM integrity. Due to the envelope features of Brucella, separating the IM and OM has not been possible, which hinders the identification of altered membrane components in a mapB mutant. For this reason, to understand the role of TAM in envelope biogenesis, Ochrobactrum anthropi was used as a closely related bacterial species. A mutant in the mapB homologue locus of O. anthropi was generated and envelope-related phenotypes were analyzed. This mutant showed increased sensitivity to lysozyme, SDS and vancomycin, confirming that MapB of alphaproteobacteria is required for cell envelope stability. Recent findings in E. coli suggested a possible function of TamB and other AsmA-like proteins in phospholipid (PL) translocation. To give insight into that hypothesis, we analyzed the PL composition of the IM and OM in the wt and ?mapB strains of O. anthropi. Highly enriched IM and OM fractions were successfully obtained by spheroplast preparation, French press disruption and sucrose gradient ultracentrifugation. To confirm IM and OM enrichment, NADH dehydrogenase activity and Western blotting with anti-OMPs were performed. The lipids were extracted by Bligh and Dyer method and organic fractions were analyzed by 1D and 2D NMR spectroscopy. Comparison of the lipid spectra from the IM and OM fractions of

the wt and the *mapB* mutant revealed significant differences in the relative amount and composition of PLs between both strains. These results provide evidence supporting the hypothesis that TAM plays a role in PL homeostasis in Alphaproteobacteria.

Palabras clave: Keywords: Outer membrane, Ochrobactrum anthropi, AsmA-like proteins, subcellular fractionation, phospholipid.